



## REVIEW SUMMARY


## AGRICULTURE

## Environmental impacts of genetically modified crops

Frederik Noack\*, Dennis Engist†, Josephine Gantois†, Vasundhara Gaur†, Batoule F. Hyjazie†, Ashley Larsen†, Leithen K. M'Gonigle†, Anouch Missirian†, Matin Qaim†, Risa D. Sargent†, Eduardo Souza-Rodrigues†, Claire Kremen

**INTRODUCTION:** In recent decades, genetically modified (GM) crops have been widely adopted by some of the world's leading agricultural nations, but the full extent of their environmental impacts remains largely unknown. Initial studies conducted in controlled environments provided valuable insights into the mechanisms that relate GM crops to the environment but offered an incomplete picture of their indirect environmental consequences. Adopting GM crops often leads to changes in other agricultural practices, such as pesticide use, cropping, and tillage patterns, with profound environmental implications (see figure). Furthermore, with widespread adoption, these changes become large in scale and can lead to substantial economic and ecological spillovers

through markets and ecological interactions, influencing the environmental outcomes of nonadopting regions. These spillovers can also lead to opposing outcomes in adopting and nonadopting regions. For example, increased profits from GM crop adoption can incentivize agricultural expansion and intensification in adopting countries with potentially negative environmental impacts. However, the increased global supply from these changes in adopting countries can reduce expansion and intensification in nonadopting countries through market effects with potentially positive environmental implications. The goal of our review is to synthesize the recent understanding of the environmental implications of GM crop adoption, with a particular focus on indirect effects and spatial spillovers.



**Environmental impacts of genetically modified crops.** Only two GM traits have been widely adopted: herbicide tolerance and insect resistance. Adopting crops with these traits affects crop losses, pesticide use, and other management actions, including tillage and crop diversity levels. These changes in turn affect agricultural expansion, deforestation, pollution, human health, greenhouse gas emissions, and biodiversity. The environmental impacts differ across geographic scales and GM traits, leading to both positive and negative effects.

Much of the existing literature focuses on the direct effects of GM crop adoption on agricultural outcomes and, to a lesser extent, its implications for the environment and human health. Recent studies employing causal inference methods have contributed to our understanding of the combined direct and indirect impacts of GM crop adoption through environmental management changes, including effects on yields, deforestation, biodiversity, and human health. Their findings paint a nuanced picture of GM crop adoption, with mostly positive impacts on yields and mixed effects on pesticide use, biodiversity, deforestation, and human health. These studies also find negative health effects from increased glyphosate use, possibly counteracted by reduced toxicity of insecticide applications due to the adoption of insect-resistant GM crops. However, the few studies that evaluate long-term consequences suggest that short-term benefits may decrease if pest resistance is not well managed. In addition, new results show that GM crop adoption increases deforestation locally, in contrast to previous studies that found a land-sparing global effect of GM crop adoption. Lastly, the evidence for the impacts of GM crop adoption on biodiversity is mixed. For example, a recent study found that although the overall impact of GM crops on bird diversity is small, the overall effect is composed of positive effects on insectivorous species and negative effects on plant and seed-eating species. While these studies greatly advance our understanding of the direct and indirect environmental effects of GM crops, they still do not fully assess the spillover effects on areas and regions that have not adopted GM crops.

**OUTLOOK:** To date, large-scale GM crop adoption has been limited to just two traits with different environmental and human health effects: herbicide tolerance and insect resistance. New GM crops and gene-edited versions with different traits, such as drought resistance, would likely have different environmental and human health impacts. Therefore, improving regulations and resistance management, providing incentives for the development and commercialization of new traits that align with social goals and human welfare, and gathering more comprehensive and detailed environmental data, especially on biodiversity, are critical for guiding agricultural innovations toward greater sustainability and allowing an accurate assessment of their impacts. ■

The list of author affiliations is available in the full article online.

\*Corresponding author. Email: frederik.noack@ubc.ca

†These authors contributed equally to this work.

Cite this article as F. Noack et al., *Science* 385, eado9340 (2024). DOI: 10.1126/science.ad9340

**S READ THE FULL ARTICLE AT**  
<https://doi.org/10.1126/science.ad9340>

## REVIEW

## AGRICULTURE

## Environmental impacts of genetically modified crops

Frederik Noack<sup>1\*</sup>, Dennis Engist<sup>1†</sup>, Josephine Gantois<sup>1,2†</sup>, Vasundhara Gaur<sup>3†</sup>, Batoule F. Hyjazie<sup>4†</sup>, Ashley Larsen<sup>5†</sup>, Leithen K. M'Gonigle<sup>6†</sup>, Anouch Missirian<sup>7†</sup>, Matin Qaim<sup>8,9†</sup>, Risa D. Sargent<sup>4†</sup>, Eduardo Souza-Rodrigues<sup>10,11†</sup>, Claire Kremen<sup>2</sup>

Genetically modified (GM) crops have been adopted by some of the world's leading agricultural nations, but the full extent of their environmental impact remains largely unknown. Although concerns regarding the direct environmental effects of GM crops have declined, GM crops have led to indirect changes in agricultural practices, including pesticide use, agricultural expansion, and cropping patterns, with profound environmental implications. Recent studies paint a nuanced picture of these environmental impacts, with mixed effects of GM crop adoption on biodiversity, deforestation, and human health that vary with the GM trait and geographic scale. New GM or gene-edited crops with different traits would likely have different environmental and human health impacts.

Genetically modified (GM) crops have been proposed as a solution to the dual challenge of meeting the demands of a growing global population while mitigating the environmental impacts of agriculture. By enhancing resistance to pests and other environmental stressors, GM traits have been promoted as having the potential to boost agricultural production without expanding agriculture into natural habitats or intensifying agrochemical applications (1–3).

Although concerns about direct toxic effects of GM crops on nontarget species and human health have attenuated recently (4–6), increasing evidence suggests that management changes arising from GM crop adoption, including shifts in pesticide use, spread of monocultures, and local agricultural expansion, can have profound implications for human health and the environment (7–9).

Quantifying the positive and negative indirect effects of GM crop adoption on the environment and human health is challenging for several reasons: First, results from field trials are only partially helpful for understanding the real-world implications of GM crop adoption, as they often hold other management factors

constant and are thus uninformative about the broader environmental implications of these indirect management changes. Second, large-scale changes in agriculture, including the widespread adoption of GM crops, also affect nonadopting farmers through changes in crop prices and environmental spillovers. Examples of those spillovers include changes in pest population sizes (10), pesticide drift (11), the development of pesticide-resistant pest populations (12, 13), and crop price effects (14, 15) that incentivize agricultural expansion or contraction and changes in the use of agrochemicals elsewhere (16–19). Such spillovers also present a methodological challenge for isolating the causal effects of GM crop adoption on agricultural outcomes and the environment.

Recent advances in causal inference techniques hold promise for analyzing the real-world consequences of widespread GM crop adoption. Examples include the quantification of GM crop adoption impacts on health (7, 20), deforestation (8), and biodiversity (9), although investigating spillovers and feedback through markets from large-scale adoption remains challenging.

We summarize the literature on the environmental impacts of GM crop adoption and highlight pathways to fill remaining knowledge gaps. Our review primarily examines the impacts of GM crops that have already been widely adopted, but we conclude by discussing the potential effects of GM and gene-edited crops still in development. Also, we take as the counterfactual a world without GM crops but with an otherwise similar conventional production system.

Genetically modifying crop germplasm involves using modern biotechnological methods to achieve specific design objectives. The environmental effects of GM crops vary depending on their specific traits. Although many traits have been developed to date, only two GM traits

have been widely commercialized. These traits are herbicide tolerance (HT), which makes a given crop tolerant to certain broad-spectrum herbicides, and insect resistance, in which genes from the bacterium *Bacillus thuringiensis* (Bt) make the crop resistant to lepidopteran insect pests. These traits hold significant commercial value as farmers worldwide struggle with weeds and lepidopteran insect pests such as corn borers, armyworms, and bollworms (21).

Adopting these two GM traits can reduce crop losses by enhancing weed and pest control and consequently increase crop yields and profits. It can also affect the use of chemical pesticides and other management practices, which may further enhance yields and profitability. We discuss these direct and indirect environmental implications below.

## GM trait development and regulation

For nearly three decades, GM crop adoption has been limited to two main GM traits, HT and Bt, in four crops: soybean, corn, cotton, and canola. Since the initial approval of those traits for cultivation, many countries have introduced stricter and more expensive regulations for GM crop approval. As a result, many other GM traits developed in labs are rarely approved for commercial application (22), including traits for resistance to fungi and bacteria, tolerance to drought and heat, and improved nitrogen use efficiency (23). It is estimated that the cost of regulation for a single new GM trait is more than USD 40 million for the trait developer (24). This regulatory cost is more than most public institutions and small and medium enterprises can afford (25), which not only prevents the commercialization of new GM traits but also contributes to a market dominated by a few large companies (26–28). These companies often patent their technologies and can inflate seed prices to raise their profits. Although increased profits from technology development incentivize innovation by those companies, it simultaneously limits seed production by farmers and competing firms (29). The extent of market concentration is visible in the market for GM seeds. In 2020, four companies controlled over half of the world's seed market. These companies hold intellectual property rights to 95% of cotton and corn varieties and 84% of soybean varieties (30). In addition to regulatory hurdles and market concentration, growing consumer concerns about GM foods have prevented companies and countries from commercializing GM versions of typical food crops such as wheat and rice (26, 31).

## GM crop adoption

Once legally approved, the decision to adopt GM crops by farmers hinge on the expected profits of adoption and the associated risks (32), the availability of the technology and its

<sup>1</sup>Food and Resource Economics, University of British Columbia, Vancouver, BC, Canada. <sup>2</sup>Institute for Resources, Environment and Sustainability, University of British Columbia, Vancouver, BC, Canada. <sup>3</sup>Institute for Policy Integrity, New York University School of Law, New York, NY. <sup>4</sup>Land and Food Systems, University of British Columbia, Vancouver, BC, Canada. <sup>5</sup>Bren School of Environmental Science & Management, University of California, Santa Barbara, Santa Barbara, CA. <sup>6</sup>Department of Biological Sciences, Simon Fraser University, Burnaby, BC, Canada. <sup>7</sup>Toulouse School of Economics, INRAe, University of Toulouse Capitole, Toulouse, France. <sup>8</sup>Center for Development Research (ZEF), University of Bonn, Germany. <sup>9</sup>Institute for Food and Resource Economics, University of Bonn, Germany. <sup>10</sup>Department of Economics, University of Toronto, Toronto, Ontario, Canada. <sup>11</sup>Centre for Economic Policy Research, London, England.

\*Corresponding author. Email: frederik.noack@ubc.ca

†These authors contributed equally to this work.

alternatives, marketing exposure (33), and farm characteristics including farm size and farmer education level (34). Recent work demonstrates that herbicide drift that damages neighboring non-GM crops can further accelerate the adoption process (17).

Following their initial commercialization in 1996, GM varieties of corn, cotton, soybean, and canola saw swift adoption rates (Fig. 1). By 2019, these GM varieties expanded to 190 million hectares representing 13% of global arable land and were cultivated in 29 countries across various income levels (Fig. 2). Among the 14 GM crops approved for cultivation, four dominate nearly 99% of the GM cultivation area, including GM soybeans (48.2% of the total GM area), GM corn (32%), cotton (13.5%), and canola (5.3%). While the HT trait is predominant in GM soybeans and canola, the Bt trait is common in GM cotton and corn, with some varieties in the USA and other nations featuring both, as “stacked” traits (35).

More than half of the global GM crop area is concentrated in five countries: the USA (38% of the total global GM crop area), Brazil (28%), Argentina (13%), Canada (7%), and India (6%). Strict regulation prevents the adoption of GM crops in many non-adopting countries. An example of the impact of regulations occurred in Romania, which grew HT soybeans between 1999 and 2006 (with an adoption rate of 80% by the end of 2006) but had to suspend cultivation upon joining the European Union (EU) in 2007 (36).

Although strong patent protection and high seed prices could limit GM crop adoption they have not been major constraints, primarily because protection of intellectual property rights is not uniform across countries. The International Union for the Protection of New Varieties of Plants (UPOV, 1961) and the World

Trade Organization’s Agreement on Trade-Related Aspects of Intellectual Property Rights (TRIPS, 1994) are the two main international agreements that govern intellectual property rights around innovation in plant and seed varieties. However, several countries are not signatories to the UPOV agreement and are therefore not required to apply its standards. Additionally, the TRIPS Agreement has provisions in Article 27(3)(b) that allow member nations to develop their own patents or other forms of protection (37, 38). Consequently, in many low- and middle-income countries, GM crops are not patented or patent protection is not strictly enforced when it does exist. Once GM crop technology receives approval in these countries, seed production and distribution face fewer legal barriers (39). This legal landscape contributes to lower seed prices in many low- and middle-income countries. Examples include Bt cotton in China, India, and Pakistan (23). In addition, some middle-income countries have begun developing generic GM varieties to facilitate small-scale farmers’ access to GM crops. Notably, Indian universities have released three cost-effective, reusable Bt cotton varieties (40), and a public-private partnership in Bangladesh has successfully introduced a Bt eggplant variety embraced by small-scale farmers (41).

#### Quantifying the causal effects of GM crops on agricultural and environmental outcomes

Despite numerous studies analyzing the effects of GM crops on yields, pesticide use, and other outcomes, several challenges persist in accurately evaluating the impacts of this technology: First, although experimental setups comparing GM crops with their non-GM counterparts can isolate the causal impact of GM technology on outcomes by holding all


other inputs fixed, these may not reflect real-world outcomes. This discrepancy occurs because farmers who adopt GM crops might also alter other agricultural practices, such as pesticide and fertilizer use, which also affect agricultural and environmental outcomes (14).

Secondly, in more realistic, nonexperimental settings, comparing outcomes between farmers (or countries) who adopt GM crops and those who do not can be confounded as adopters and non-adopters often differ in attributes such as farm size, education level, access to irrigation, and labor constraints (34), which can influence the outcomes of interest independently of GM crop adoption and related agricultural practices. Some of these differences can be accounted for with proper statistical techniques [e.g., (42)]. A global meta-analysis of the many existing studies with non-experimental farm-level data suggests that GM crop adoption contributes to significant yield gains and that the Bt trait leads to reductions in chemical pesticide use, whereas the HT trait does not (43). However, challenges with causal inference persist as researchers cannot observe all confounding characteristics that could potentially bias the estimates.

One approach to addressing these challenges is implementing randomized control trials (RCTs) in field settings, wherein randomly selected farmers receive incentives to adopt GM crops. These behavioral experiments enable farmers to modify their agricultural practices in conjunction with GM adoption, resulting in directly comparable groups of adopters and non-adopters. Ahmed *et al.* (41) conducted such an RCT in Bangladesh, and the results of this study showed a 50% increase in yields and a 40% reduction in pesticide costs for Bt eggplant production compared with conventional eggplant production in the control villages.


Despite RCTs being the gold standard for causal inference, they are not without limitations. First, the results from RCTs are not always scalable to larger, aggregated levels as market responses and spillovers are largely absent at the small scale of the experiments (44, 45). Secondly, because of varying local conditions these results may not be directly transferable to different settings such as other countries or types of GM crops.

Causal inference techniques that evaluate the consequences of large-scale adoption across several countries are instrumental for understanding the real-world implications of GM crop adoption. A recent study by Hansen and Wingender (46) uses one such causal inference method to estimate the impact of GM crop adoption on yields on a global scale. The study distinguished the effects of GM adoption from broader country-specific agricultural development by comparing yield differences across corn, cotton, soybean, and other crops before and after GM adoption across countries with



**Fig. 1. GM crop adoption over time.** The percentage of global corn, cotton, soybean, and canola area under GM varieties. The data are from the status reports of the ISAAA [e.g., (35)].





**Fig. 2. GM crop adoption across countries.** Percentage of global corn, cotton, soybean, and canola cropland area planted with GM varieties in 2018. Countries with zero or less than 1% GM crop adoption are shown in light gray. The data are from the status reports of the ISAAA [e.g., (35)].

and without GM crops, employing a triple difference methodology. They report no yield effects of GM soybean and rapeseed adoption but large and statistically significant yield effects of GM cotton and corn adoption. These results may overestimate the actual yield gains, however, because yield gains of such magnitude in some of the world's largest crop producers would change global crop prices and thus affect non-adopters' incentives for production. Indeed, Barrows *et al.* (14) found that GM crop adoption reduced prices between 10 and 20% compared with the counterfactual world without GM crop adoption. Although such a price reduction represents a benefit of GM crops for consumers, lower prices can diminish the revenues of non-adopting farmers, potentially prompting them to reduce land, fertilizer, and other inputs to agricultural production (16–19). To illustrate this methodological challenge, consider yield trends in adopting and non-adopting countries (Fig. 3). Corn, cotton, and soybean yields were different in adopting and non-adopting countries even before the commercialization of GM crops and generally increased over time. Yields further diverged between adopting and non-adopting countries after the commercialization of GM crops, suggesting a positive impact of GM crops on yields (Fig. 3). Causal inference approaches address the challenges arising from different yield levels before adoption and general time trends. However, if yield trends in non-adopting countries declined in response to GM commercialization due to, for example, price effects, it would also contribute to the divergence of yields between

adopting and non-adopting countries and thus lead to an overestimation of the positive yield effects of GM crops.

#### Agricultural expansion and deforestation


Yield increases from GM crop adoption may reduce the incentives to convert forests to croplands, thus reducing biodiversity loss and greenhouse gas emissions associated with land use change (15, 47). The suggested mechanism for this land-sparing effect is that increased production in response to GM crop adoption leads to reduced crop prices, discouraging farmers from expanding their cropland into natural or seminatural habitats. This general positive relationship between crop prices and deforestation, which drives the effect on agricultural expansion or contraction, has been established in various studies and locations (18, 48–50) although unrelated to GM crops.

Alternatively, increased profits from GM crop adoption, e.g., through reduced production costs or crop losses, could motivate farmers to expand agricultural production into natural habitats. For instance, Carreira *et al.* (8) find that deforestation in Brazil increased in areas with significant productivity gains from GM crop adoption compared with areas with low benefits of GM crop adoption, indicating a potential deforestation-inducing effect of GM crops.

Although research specifically linking GM crop adoption to deforestation is limited, broader studies on agricultural productivity and deforestation present varied results (51, 52). A possible explanation for these mixed results

is that the contrasting outcomes may occur simultaneously but at different scales or in different locations. For example, although GM crop adoption might lead to agricultural expansion in adopting regions as a result of increased profits, it could simultaneously reduce agricultural expansion in non-adopting regions because of the lack of profitability gains in those regions and lower crop prices in response to the increased aggregate supply. Across all regions, this could lead to overall forest loss or forest gain, depending on the magnitude of deforestation in adopting and non-adopting regions. Further, the effect depends on the response of prices to aggregate supply. For example, the land-sparing effect may be stronger in relatively closed economies with highly responsive prices whereas exporting countries facing world prices may not experience the same land-sparing effect through the price mechanism.

Although it is conceptually clear that increased crop production in one region can reduce agricultural expansion elsewhere, it is difficult to empirically quantify where cropland would be reduced and by how much. Computable general equilibrium (CGE) models are often used to quantify these spatial relationships [e.g., (53)]. For example, (54) found that the Green Revolution, characterized by the adoption of high-yielding crop varieties in low-income countries since the late 1960s, potentially prevented the conversion of 18 to 27 million hectares of non-natural vegetation into farmland, thanks to localized production increases and subsequent price effects. However, caution is warranted



**Fig. 3. GM crop adoption and crop yields.** The red lines are mean crop yields for countries with at least 10% adoption of GM corn, cotton, and soybeans in 2018, respectively. The blue lines are the mean crop yields for the other countries. The dotted line is the year of GM commercialization (1996), after which adopting countries started to increase the share of land under GM varieties. The GM adoption data are from the status reports of the ISAAA [e.g., (35)]. The yield data are from the FAO (130).

when using CGE models as they rely on numerous parametrized relationships that may introduce biases in the analysis and it often remains unclear which land cover the expanding agriculture displaces.

### Pesticide use

The two main types of GM crops, Bt and HT, are both related to pesticide use but in different ways. The Bt trait provides resistance to lepidopteran insect pests, thus reducing the need for chemical insecticide sprays against these particular pests. Numerous studies have found evidence that Bt crop adoption is associated with significant insecticide reductions both in low-income and high-income countries (41, 55–57). However, the influence of Bt crops on insecticide application may fluctuate over time depending on various factors, including management practices of farmers. First, Bt targets specific insect pests, often lepidopteran, in cotton (e.g., bollworms) and lepidopteran or coleopteran in corn (e.g., corn borers, corn rootworm). Thus, as with other pesticides, nontarget, secondary insect pests may proliferate, which could lead to a rebound effect in insecticide use (58). Second, pest populations can develop resistance to Bt, especially when

farmers do not plant refuge areas with non-Bt crops (57, 59–61) or repeatedly plant the same variety of Bt crop. For example, field-evolved resistance has been observed in corn rootworm, one of the most serious corn pests in the US (62). As with other pesticides, resistance to Bt can then lead to increased pesticide use. Evaluation of long-term data suggests that Bt technologies can remain effective for many years and that pest resistance buildup can be managed with agronomic and breeding strategies (63) such as crop rotations and diversifying the type of Bt planting (64).

By contrast, HT crops are designed to be tolerant to certain broad-spectrum herbicides (historically and most prominently, glyphosate) and are intended to be used in conjunction with herbicides. Thus, empirical studies report either no significant reduction in overall pesticide use following HT adoption (32) or a substantial increase in the quantity of herbicides used (43). Furthermore, HT crops can cause a substitution effect in which the broad-spectrum herbicides they tolerate, such as glyphosate, replace other specific herbicides used in conventional agriculture that might be more or less toxic (65). This substitution has led to a substantial increase in glyphosate

usage since the mid-1990s, particularly in countries that have widely adopted HT crops (66).

Another indirect consequence of GM crop adoption is that a reduction in the diversity of crop types and herbicide-active ingredients has contributed to herbicide resistance development in weeds (66). New GM crop varieties with multiple resistances against herbicides, such as the addition (stacking) of a dicamba-tolerance trait, have been commercialized to control these weeds. The resulting increased use of this potentially more toxic and volatile herbicide raises new concerns regarding human health (acute and chronic conditions related to dicamba), damages to non-dicamba-tolerant crops on neighboring fields (17), and harm to nontarget vegetation in surrounding ecosystems. We discuss further implications of pesticide changes for pollution, human health, and biodiversity in more detail below.

### Health effects from pesticide use

Analogous to the implications for agricultural expansion and deforestation, understanding the impacts of GM crop adoption on human health and environmental pollution necessitates examining both direct and indirect effects. Direct adverse effects on human health

from GM crop consumption are now broadly considered to be negligible (6). The indirect consequences, especially those related to changes in farmers' pesticide use, are proving significant and are receiving increasing attention.

Pesticide exposure occurs through dermal contact, ingestion, or inhalation. Farmers, farmworkers, and those involved in pesticide production are likely to experience the greatest health impacts from GM-driven changes in pesticide use (67). Populations residing near agricultural fields and the general public are exposed to pesticides primarily through diet or air and water pesticide contamination. Thus, regional pesticide regulation and residue monitoring are important indicators (68). Pesticide regulations and guidelines vary worldwide, with low-income countries often lacking the resources to implement or enforce legislation to avoid elevated occupational pesticide risks or adequately limit pesticide residues from the food supply chain (68). Thus, the health impacts of GM-driven changes in pesticides are likely to be magnified in low-income countries for both occupational and nonoccupational communities.

Following the adoption of Bt crops, reductions in insecticide use have generally led to positive environmental and human health outcomes for farmers, farm workers, and consumers (56, 69). For example, Kouser and Qaim (70), utilizing detailed panel data from India, demonstrated that Bt cotton adoption has prevented millions of insecticide poisoning incidents annually among smallholder farmers through reduced spraying. These health benefits, however, may wane over time if insecticide use on Bt crops increases as a result of the emergence of Bt-resistant pests or the proliferation of pests unaffected by Bt technology (see above).

Quantifying the health and pollution effects of HT crop adoption is complex due to changes in both the volumes and types of herbicides used (i.e., substitution of broad-spectrum herbicides for more specific ones, especially glyphosate). Nelson and Bullock (71) argue that changes in herbicide use led to a decrease in health hazards on the basis of acute toxicity metrics whereas the International Agency for Research on Cancer (IARC) (72) classifies glyphosate as a probable carcinogen and suggests a net increase in cancer risk (73). Beyond the active ingredients of the herbicides, adjuvants (compounds that enhance the spread and effectiveness of herbicides) also contribute to the overall impact of pesticide application. For example, common adjuvants used with glyphosate, such as polyethoxylated tallow amine (POEA), have been shown to be substantially more toxic to humans (74) than glyphosate alone, and adversely affect aquatic and terrestrial species (12, 75), a point that we discuss further below.

The risk quotient, a common metric for pesticide toxicity that combines pesticide quantities with health risk factors (e.g., measured as the inverse of the amount lethal to half of a rat population), offers insight into the health effects of pesticides (76, 77). A study by Lee *et al.* (78) applied this metric and found that GM crop adoption in the USA initially reduced pesticide toxicity. Over time, however, toxicity levels rose, partly due to the development of pesticide resistance. Overall, it is important to note that ex-ante toxicity studies might be subject to three main biases: (i) pesticide toxicity measurements can be imprecise (79, 80), (ii) evaluation may focus only on active ingredients ignoring the toxicity of adjuvants (74, 75, 81), and (iii) the studies may measure hazard (i.e., potential of harm; toxicity) instead of risk (i.e., probability of harm; the interaction of toxicity with exposure), potentially under- or overestimating the actual damage to humans and nontarget species (82, 83).

An alternative approach to assessing the health impacts of GM crops involves directly measuring health outcomes associated with GM crop adoption through causal inference methods. These methods leverage natural variations, such as differences in GM crop suitability across regions or variations in the distribution of environmental impacts (e.g., upstream and downstream of GM crop adoption), to establish the causal effects of GM crops on health. These methods were used in Brazil, where widespread adoption of HT soybeans led to increased glyphosate use (7, 84). Dias *et al.* (7) found that HT crop adoption and the resulting rise in glyphosate levels in water significantly increased infant mortality, pre-term birth rates, and the occurrence of low birth weights. Similarly, Skidmore *et al.* (84) linked the heightened pesticide use to elevated childhood cancer rates, likely through water contamination. Rubin and Saulnier (20) show similar negative health outcomes in response to increased glyphosate use from GM crop adoption in the US.

Although significant progress has been made in assessing the environmental and health impacts of pesticide changes due to GM crop adoption, considerable uncertainties remain. These uncertainties relate to the effects of changes in the type, amount, and application timing of pesticides used on GM fields.

#### Further management practices

GM crop adoption has been associated with changes in other management practices beyond pesticide use, though the evidence is sparse, and the overall environmental consequences of these changes are still largely unknown. First, GM crops—especially those tolerant to glyphosate—may have facilitated reduced tillage intensity, achieved through conservation tillage or no-till practices, due to more effective weed control

(85–87). For example, Perry *et al.* (88) observe that glyphosate-tolerant soybeans have promoted the use of these practices, increasing conservation tillage and no-till by approximately 10 and 20%, respectively. Although the impact of reduced tillage practices on productivity is mixed (89), they have been shown to benefit soil structure, runoff, water quality, soil biota, aboveground wildlife, and air quality with varying levels of magnitude (90–92); reduced tillage practices also have mixed impacts on carbon sequestration (93, 94) (see below).

Second, GM crop adoption may also influence crop rotation practices. Specifically, GM crops, by providing effective weed and pest control, might reduce the traditional yield advantages gained from rotating crops (95). This substitution effect could potentially reduce crop rotations at the field level and lead to more crop uniformity across larger agricultural landscapes, thereby reducing both temporal and spatial crop diversity. Causal evidence for this relationship has yet to be established. Understanding this relationship is crucial, as diverse crop rotations support wild biodiversity (96), reduce pesticide applications in the agricultural landscape (97), and enhance the resilience of agricultural systems to adverse weather conditions (98).

Finally, new crop varieties can encourage increased use of further inputs such as fertilizer (99), which can have profoundly harmful downstream consequences for the environment and human health (100). Below, we discuss the implications of these management changes for biodiversity and greenhouse gas emissions.

#### Biodiversity

GM crop adoption can affect biodiversity directly through the consumption of GM crops or indirectly through various land use changes, including agricultural expansion, pesticide use, and other management practices. Considerable literature has examined the direct effects of GM crops on nontarget species. Most studies focus on the effects of Bt crop consumption on a diversity of organisms; these studies have identified adverse toxic effects on a range of nontarget species, including butterflies, springtails, and lacewings (4, 101–103). Although these studies highlight potential risks, they also underscore substantial gaps in our knowledge, particularly regarding aquatic, microbial, and soil-dwelling organisms (104, 105). In addition, the concern has been raised that many studies are poorly replicated, conducted in nonfield realistic conditions, and often funded by industry stakeholders (4, 101). Given these caveats, the available evidence suggests that, for the species tested, risks posed by direct and immediate contact with GM crops under field-realistic conditions appear small or nonexistent (5).

The movement of transgenes into wild populations is another oft-cited risk of GM

crop adoption (106). Gene flow from GM crops to wild or locally adapted crops could reduce biological diversity through genetic assimilation (107) or give rise to new lineages that persist (108). There is considerable evidence of gene flow from GM to wild species, for example, in the case of GM to wild canola varieties in Canada (14). At the same time, the fate of lineages resulting from transgenic gene flow is inherently difficult to predict (109) and, to date, the risk of negative environmental or economic impact appears to be low (110).

GM crop adoption has led to several changes in global pesticide usage (see above) and there is a clear scientific consensus that pesticides threaten biodiversity worldwide (111–114). In addition to the direct impacts of Bt and herbicides on wildlife living within GM crops, herbicide spray drift can lead to the loss of wild plant diversity in nearby landscapes, even at low concentrations (115). This loss of host plant diversity can impose negative downstream effects on key ecosystem players such as pollinators and other nontarget organisms (116).

Finally, there is growing consensus that changes in agricultural management practices that reduce crop and landscape heterogeneity (i.e., agricultural simplification) also negatively affect biodiversity. For instance, Strobl (96) found that greater crop diversity is linked to higher bird diversity in agricultural landscapes. Furthermore, a comprehensive meta analysis by Sirami *et al.* (117) provided evidence that agricultural landscape simplification is associated with lower multitrophic biodiversity. Agricultural expansion, when it occurs at the expense of natural and seminatural habitats, is the single largest cause of biodiversity reduction through wildlife habitat loss and degradation (118, 119). Both types of land use change—agricultural simplification and expansion into natural areas—have been linked to GM crop adoption (see above), although the exact extent of GM crops' role in simplification and expansion remains uncertain. A recent study that evaluated the combined direct and indirect effects of GM crop adoption on bird diversity in the US found overall positive effects of GM crop adoption on bird diversity with heterogeneous effects across species groups and crops. Although insectivorous birds benefited from GM crop adoption, especially from GM cotton, herbivorous birds declined in response to GM crop adoption, with GM soy having the strongest impact (9).

A major challenge in evaluating the risks associated with GM crops for biodiversity is the scarcity of standardized, long-term biodiversity data from agricultural landscapes. Although reliable longitudinal bird survey data exist in some countries, such comprehensive data are lacking for other critical taxa, including insects, microbes, and plants. The absence of long-term biodiversity records for taxonomic

groups heavily affected by pesticides, such as insects and microbes, hinders a full understanding of GM crop management's effects on biodiversity (120–123). These groups, which also play crucial functional and trophic roles (e.g., as mutualists, prey, predators, and pathogens) in the same agroenvironments, require more extensive research to fully grasp the broader impacts of GM crop adoption on biodiversity, particularly taxa with limited existing survey records.

### Greenhouse gas emissions and climate change

GM crop adoption can affect greenhouse gas emissions through agricultural expansion and deforestation and changes in management practices on existing agricultural land, including tillage practices, fertilizer applications, the use of agricultural machinery, and shifts between crop and livestock production (47, 124). Collectively, these changes could significantly affect greenhouse gas emissions (15). For instance, Kovak *et al.* (47) estimated that GM crop adoption in Europe could reduce agricultural greenhouse gas emissions by 7.5% as a result of higher yields and thus reduce agricultural expansion at the global level. In terms of tillage, adopting GM crops with the HT trait has facilitated no-till practices, reducing tractor and fuel use and possibly increasing soil carbon sequestration. However, the benefits of no-till agriculture for reducing greenhouse gas emissions are controversial and may require integration with other practices, including cover cropping and crop rotations (89, 94). Further changes in greenhouse gas emissions can result from fertilizer adjustments (see above), but the impact of GM crop adoption on emissions from fertilizer use is largely unknown.

The prevailing view in the current literature is that GM crop adoption could be greenhouse gas-reducing (47, 124), but substantial uncertainties over the magnitudes of greenhouse gas emissions still persist. So far, most GM crop applications are found in conventionally managed systems with relatively high greenhouse gas emissions per unit of land.

### Outlook

GM crop adoption has been rapid and complete throughout several of the major crop-producing countries in the world. However, this adoption is mostly confined to two GM traits (HT and Bt) and to a small number of commercial crops. Although some countries have embraced these GM technologies, others, including members of the European Union and countries in Africa and Asia, still ban their production mainly because of perceived environmental and health concerns.

Much progress has been made in recent years to quantify the impact of GM crop adoption on agricultural inputs. These studies are fundamental for understanding the pathways through which GM crop adoption could

affect environmental and health outcomes. However, linking those individual management changes to the relevant outcomes is complicated by the number of pathways that could affect the outcomes. Examples include the indirect impacts of GM crops on biodiversity through changes in land use and cover, pesticide toxicity, or species interactions. Focusing on individual impacts, such as pesticide toxicity, may lead to an incomplete picture of the overall effects. A growing literature, therefore, focuses on the combined indirect and direct effects of GM crop adoption on environmental and health outcomes using causal inference techniques and “natural” experiments. These studies include biodiversity (9), health (7), and forest cover (8).

Despite substantial progress, large knowledge gaps remain. First, large uncertainties persist regarding the long-term impact of GM crops on the expansion of monocultures and the spread of resistant weeds and pests. These changes may diminish or even reverse the short-term benefits of pesticide reductions from GM crop adoption (78). Second, the long-term impact of GM crop adoption on species groups other than birds, including bees, butterflies, and other insects, is largely unassessed. These groups may be directly affected by GM crops and pesticide use changes, and their abundance and diversity may in turn directly affect agricultural production (125). The lack of systematic long-term surveys of these species groups hampers progress in this field. Third, the quantification of GM crop adoption on deforestation on a global scale has not been quantified. Although local deforestation effects have been demonstrated, e.g., in Brazil where deforestation increased in response to GM crop adoption (8), the often-assumed reduction of deforestation at a global scale in response to GM crop adoption has not been tested empirically.

It is important to note that the environmental consequences of GM crop adoption evaluated here have largely been assessed in similar farming systems, namely simplified conventional monoculture systems, setting a low bar for environmental comparisons. Such simplified systems are known to substantially reduce both biodiversity and ecosystem services relative to more diversified forms of agriculture (126–128). Thus, any positive environmental effects of GM crops, such as improvements to bird biodiversity (9), must be interpreted cautiously. While GM crops that maintain or increase yields while reducing environmental harm may be developed, it is critical to go beyond harm reduction toward regenerating environmental benefits to achieve agricultural sustainability.

Although this review finds mixed effects of current GM crops on the environment, it is important to remember that current GM crop production is dominated by a few GM-crop-trait



combinations that were selected in a process driven by industry self-interest, public acceptance (or lack thereof), and costly regulation. It is plausible that GM crops with other traits explicitly developed to reduce the environmental impacts of agriculture combined with stringent resistance management efforts would have unambiguous positive environmental consequences. New gene-editing tools can reduce the cost and increase the speed and precision of developing desirable traits. However, this will require a conducive policy and regulatory environment that fosters diversity, transparency, sustainability, and a less concentrated industry (33, 129). Finally, GM and gene-edited crops and traits should not be seen as a substitute for good agronomic practices but should be integrated smartly into sustainable production systems.

## REFERENCES AND NOTES

- L. T. Hickey *et al.*, Breeding crops to feed 10 billion. *Nat. Biotechnol.* **37**, 744–754 (2019). doi: [10.1038/s41587-019-0152-9](https://doi.org/10.1038/s41587-019-0152-9); pmid: [31209375](https://pubmed.ncbi.nlm.nih.gov/31209375/)
- S. S. Zaidi *et al.*, New plant breeding technologies for food security. *Science* **363**, 1390–1391 (2019). doi: [10.1126/science.aav6316](https://doi.org/10.1126/science.aav6316); pmid: [30923209](https://pubmed.ncbi.nlm.nih.gov/30923209/)
- J. Hazel, J. Jones, “Enabling Genetic Technologies for Food Security” (The Royal Society, 2023).
- M. Marvier, C. McCreedy, J. Regetz, P. Kareiva, A meta-analysis of effects of Bt cotton and maize on nontarget invertebrates. *Science* **316**, 1475–1477 (2007). doi: [10.1126/science.1139208](https://doi.org/10.1126/science.1139208); pmid: [17556584](https://pubmed.ncbi.nlm.nih.gov/17556584/)
- M. Meissle, S. E. Naranjo, J. Romeis, Does the growing of Bt maize change abundance or ecological function of non-target animals compared to the growing of non-GM maize? A systematic review. *Environ. Evid.* **11**, 21 (2022). doi: [10.1186/s13750-022-00272-0](https://doi.org/10.1186/s13750-022-00272-0)
- Genetically Engineered Crops: Experiences and Prospects* (The National Academies Press, 2016).
- M. Dias, R. Rocha, R. R. Soares, Down the River: Glyphosate Use in Agriculture and Birth Outcomes of Surrounding Populations. *Rev. Econ. Stud.* **90**, 2943–2981 (2023). doi: [10.1093/restud/rdad011](https://doi.org/10.1093/restud/rdad011)
- I. Carreira, F. Costa, J. P. Pessoa, The deforestation effects of trade and agricultural productivity in Brazil. *J. Dev. Econ.* **167**, 103217 (2024). doi: [10.1016/j.jdevco.2023.103217](https://doi.org/10.1016/j.jdevco.2023.103217)
- D. Engist, M. Guzman, A. Larsen, T. Church, F. Noack, The impact of genetically modified crops on bird diversity. *Nat. Sustain.* (2024). doi: [10.1038/s41893-024-01390-y](https://doi.org/10.1038/s41893-024-01390-y)
- W. D. Hutchison *et al.*, Areawide suppression of European corn borer with Bt maize reaps savings to non-Bt maize growers. *Science* **330**, 222–225 (2010). doi: [10.1126/science.1190242](https://doi.org/10.1126/science.1190242); pmid: [20929774](https://pubmed.ncbi.nlm.nih.gov/20929774/)
- A. Missirian, “Yes, in your backyard: Forced technological adoption and spatial externalities” (TSE Working Papers, 2024).
- G. Schütte *et al.*, Herbicide resistance and biodiversity: Agronomic and environmental aspects of genetically modified herbicide-resistant plants. *Environ. Sci. Eur.* **29**, 5 (2017). doi: [10.1186/s12302-016-0100-y](https://doi.org/10.1186/s12302-016-0100-y); pmid: [28163993](https://pubmed.ncbi.nlm.nih.gov/28163993/)
- A. R. Kniss, Genetically engineered herbicide-resistant crops and herbicide-resistant weed evolution in the United States. *Weed Sci.* **66**, 260–273 (2018). doi: [10.1017/wsc.2017.70](https://doi.org/10.1017/wsc.2017.70)
- G. Barrows, S. Sexton, D. Zilberman, Agricultural biotechnology: The promise and prospects of genetically modified crops. *J. Econ. Perspect.* **28**, 99–120 (2014). doi: [10.1257/jep.28.1.99](https://doi.org/10.1257/jep.28.1.99)
- G. Barrows, S. Sexton, D. Zilberman, The impact of agricultural biotechnology on supply and land-use. *Environ. Dev. Econ.* **19**, 676–703 (2014). doi: [10.1017/S1355770X14000400](https://doi.org/10.1017/S1355770X14000400)
- J. Hargrave, K. Kis-Katos, Economic causes of deforestation in the Brazilian Amazon: A panel data analysis for the 2000s. *Environ. Resour. Econ.* **54**, 471–494 (2013). doi: [10.1007/s10640-012-9610-2](https://doi.org/10.1007/s10640-012-9610-2)
- J. Fernandez-Cornejo, R. F. Nehring, C. Osteen, S. Wechsler, A. Martin, A. Vialou, “Pesticide Use in U.S. Agriculture: 21 Selected Crops, 1960–2008” (US Department of Agriculture, technical report EIB-124, 2014).
- J. Assunção, C. Gaudour, R. Rocha, Deforestation slowdown in the Brazilian Amazon: Prices or policies? *Environ. Dev. Econ.* **20**, 697–722 (2015). doi: [10.1017/S1355770X15000078](https://doi.org/10.1017/S1355770X15000078)
- R. Miao, M. Khanna, H. Huang, Responsiveness of crop yield and acreage to prices and climate. *Am. J. Agric. Econ.* **98**, 191–211 (2016). doi: [10.1093/ajae/aav025](https://doi.org/10.1093/ajae/aav025)
- E. Rubin, E. Saulnier, “Glyphosate exposure and GM seed rollout unequally reduced perinatal health” (SSRN, 2024); [https://papers.ssrn.com/sol3/papers.cfm?abstract\\_id=4867914](https://papers.ssrn.com/sol3/papers.cfm?abstract_id=4867914)
- S. Savary *et al.*, The global burden of pathogens and pests on major food crops. *Nat. Ecol. Evol.* **3**, 430–439 (2019). doi: [10.1038/s41559-018-0793-y](https://doi.org/10.1038/s41559-018-0793-y); pmid: [30718852](https://pubmed.ncbi.nlm.nih.gov/30718852/)
- J. Wesseler, R. D. Smart, J. Thomson, D. Zilberman, Foregone benefits of important food crop improvements in Sub-Saharan Africa. *PLOS ONE* **12**, e0181353 (2017). doi: [10.1371/journal.pone.0181353](https://doi.org/10.1371/journal.pone.0181353); pmid: [28749984](https://pubmed.ncbi.nlm.nih.gov/28749984/)
- M. Qaim, Role of new plant breeding technologies for food security and sustainable agricultural development. *Appl. Econ. Perspect. Policy* **42**, 129–150 (2020). doi: [10.1002/aep.13044](https://doi.org/10.1002/aep.13044)
- Agbiolinvestor, “Time and cost to develop a new GM trait” (Agbiolinvestor, technical report, 2022).
- B. Schiek *et al.*, Demystification of GM crop costs: Releasing late blight resistant potato varieties as public goods in developing countries. *Int. J. Biotechnol.* **14**, 112–131 (2016). doi: [10.1504/IJBT.2016.077942](https://doi.org/10.1504/IJBT.2016.077942)
- R. Herring, R. Paarlberg, The political economy of biotechnology. *Annu. Rev. Resour. Econ.* **8**, 397–416 (2016). doi: [10.1146/annurev-resource-100815-095506](https://doi.org/10.1146/annurev-resource-100815-095506)
- M. F. Brennan, C. E. Pray, A. Courtmanche, “Impact of industry concentration on innovation in the US plant biotech industry,” in *Proceedings of NE-165 Conference* (2000), pp. 153–174.
- P. H. Howard, Visualizing consolidation in the global seed industry: 1996–2008. *Sustainability* **1**, 1266–1287 (2009). doi: [10.3390/sui041266](https://doi.org/10.3390/sui041266)
- M. Qaim, A. De Janvry, Genetically modified crops, corporate pricing strategies, and farmers’ adoption: The case of Bt cotton in Argentina. *Am. J. Agric. Econ.* **85**, 814–828 (2003). doi: [10.1111/1467-8276.00490](https://doi.org/10.1111/1467-8276.00490)
- USDA Agricultural Marketing Service, “More and Better Choices for Farmers: Promoting Fair Competition and Innovation in Seeds and Other Agricultural Inputs” (USDA, 2023).
- R. Paarlberg *et al.*, The uptake of new crop science: Explaining success, and failure. *Food Policy* **122**, 102572 (2024). doi: [10.1016/j.foodpol.2023.102572](https://doi.org/10.1016/j.foodpol.2023.102572)
- J. Fernandez-Cornejo, S. J. Wechsler, M. Livingston, L. Mitchell, “Genetically Engineered Crops in the United States” (USDA, Economic Research Service, technical report ERR-162, 2014).
- J. Clapp, The problem with growing corporate concentration and power in the global food system. *Nat. Food* **2**, 404–408 (2021). doi: [10.1038/s43016-021-00297-7](https://doi.org/10.1038/s43016-021-00297-7); pmid: [37118223](https://pubmed.ncbi.nlm.nih.gov/37118223/)
- U. Aldana, B. Barham, J. Foltz, P. Useche, Early adoption, experience, and farm performance of GM corn seeds. *Agric. Econ.* **43**, 11–18 (2012). doi: [10.1111/j.1574-0862.2012.00616.x](https://doi.org/10.1111/j.1574-0862.2012.00616.x)
- ISAAA, “Global status of commercialized biotech/GM crops in 2019” (International Service for the Acquisition of Agri-biotech Applications, 2019).
- C. James, ISAAA, “Global status of commercialized biotech/GM crops: 2007” (ISAAA, brief 37, 2007).
- B. Dhar, Quaker International Nations Office (QUONO), “Sui generis systems for plant variety protection” (QUONO, 2002).
- D. Leskien, M. Flitner, “Intellectual property rights and plant genetic resources: Options for a sui generis system” (International Plant Genetic Resources Institute, 1997).
- S. Boettiger, G. Graff, P. G. Pardey, E. Van Dusen, B. D. Wright, “Intellectual property rights for plant biotechnology: International aspects” in *Handbook of Plant Biotechnology*, Ch. 56, pp. 1088–1113 (2004).
- ISAAA, “Indian Universities Develop Bt Cotton Varieties with Reusable Seeds” (ISAAA, 2017); <https://www.isaaa.org/kc/croppbiotechupdate/article/default.asp?ID=15257>
- A. U. Ahmed, J. Hoddinott, N. Abedin, N. Hossain, The impacts of GM foods: Results from a randomized controlled trial of Bt eggplant in Bangladesh. *Am. J. Agric. Econ.* **103**, 1186–1206 (2021). doi: [10.1111/ajae.12162](https://doi.org/10.1111/ajae.12162)
- J. Kathage, M. Qaim, Economic impacts and impact dynamics of Bt (*Bacillus thuringiensis*) cotton in India. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 11652–11656 (2012). doi: [10.1073/pnas.1203647109](https://doi.org/10.1073/pnas.1203647109); pmid: [22753493](https://pubmed.ncbi.nlm.nih.gov/22753493/)
- W. Klümper, M. Qaim, A meta-analysis of the impacts of genetically modified crops. *PLOS ONE* **9**, e111629 (2014). doi: [10.1371/journal.pone.0111629](https://doi.org/10.1371/journal.pone.0111629); pmid: [25365303](https://pubmed.ncbi.nlm.nih.gov/25365303/)
- K. Muralidharan, P. Niehaus, Experimentation at scale. *J. Econ. Perspect.* **31**, 103–124 (2017). doi: [10.1257/jep.31.4.103](https://doi.org/10.1257/jep.31.4.103)
- C. A. Davis, A. M. Mobarak, The challenges of scaling effective interventions: A path forward for research and policy. *World Dev.* **127**, 104817 (2020). doi: [10.1016/j.worlddev.2019.104817](https://doi.org/10.1016/j.worlddev.2019.104817)
- C. W. Hansen, A. M. Wingender, *Am. Econ. Rev. Insights* **5**, 224–240 (2023). doi: [10.1257/aeri.20220144](https://doi.org/10.1257/aeri.20220144)
- E. Kovak, D. Blaustein-Rejto, M. Qaim, Genetically modified crops support climate change mitigation. *Trends Plant Sci.* **27**, 627–629 (2022). doi: [10.1016/j.tplants.2022.01.004](https://doi.org/10.1016/j.tplants.2022.01.004); pmid: [35148945](https://pubmed.ncbi.nlm.nih.gov/35148945/)
- T. Harding, J. Herzberg, K. Kuralbayeva, Commodity prices and robust environmental regulation: Evidence from deforestation in Brazil. *J. Environ. Econ. Manage.* **108**, 102452 (2021). doi: [10.1016/j.jeem.2021.102452](https://doi.org/10.1016/j.jeem.2021.102452)
- C. Balboni, A. Berman, R. Burgess, B. A. Olken, The economics of tropical deforestation. *Annu. Rev. Econ.* **15**, 723–754 (2023). doi: [10.1146/annurev-economics-090622-024705](https://doi.org/10.1146/annurev-economics-090622-024705)
- N. Berman, M. Couttenier, A. Leblois, R. Soubeyran, Crop prices and deforestation in the tropics. *J. Environ. Econ. Manage.* **119**, 102819 (2023). doi: [10.1016/j.jeem.2023.102819](https://doi.org/10.1016/j.jeem.2023.102819)
- R. Abman, C. Carney, Agricultural productivity and deforestation: Evidence from input subsidies and ethnic favoritism in Malawi. *J. Environ. Econ. Manage.* **103**, 102342 (2020). doi: [10.1016/j.jeem.2020.102342](https://doi.org/10.1016/j.jeem.2020.102342)
- B. Dalheimer, C. Kubitz, B. Brümmer, Technical efficiency and farmland expansion: Evidence from oil palm smallholders in Indonesia. *Am. J. Agric. Econ.* **104**, 1364–1387 (2022). doi: [10.1111/ajae.12267](https://doi.org/10.1111/ajae.12267)
- T. W. Hertel, N. Ramankutty, U. L. C. Baldos, Global market integration increases likelihood that a future African Green Revolution could increase crop land use and CO2 emissions. *Proc. Natl. Acad. Sci. U.S.A.* **111**, 13799–13804 (2014). doi: [10.1073/pnas.1403543111](https://doi.org/10.1073/pnas.1403543111); pmid: [25201962](https://pubmed.ncbi.nlm.nih.gov/25201962/)
- J. R. Stevenson, N. Villoria, D. Byerlee, T. Kelley, M. Maredia, Green Revolution research saved an estimated 18 to 27 million hectares from being brought into agricultural production. *Proc. Natl. Acad. Sci. U.S.A.* **110**, 8363–8368 (2013). doi: [10.1073/pnas.1208065110](https://doi.org/10.1073/pnas.1208065110); pmid: [23671086](https://pubmed.ncbi.nlm.nih.gov/23671086/)
- J. Huang *et al.*, A decade of Bt cotton in Chinese fields: Assessing the direct effects and indirect externalities of Bt cotton adoption in China. *Sci. China Life Sci.* **53**, 981–991 (2010). doi: [10.1007/s11427-010-4036-y](https://doi.org/10.1007/s11427-010-4036-y); pmid: [20821297](https://pubmed.ncbi.nlm.nih.gov/20821297/)
- P. C. Veetil, V. V. Krishna, M. Qaim, Ecosystem impacts of pesticide reductions through Bt cotton adoption. *Aust. J. Agric. Resour. Econ.* **61**, 115–134 (2017). doi: [10.1111/1467-8489.12171](https://doi.org/10.1111/1467-8489.12171)
- A. J. Gassmann, D. D. Reisig, Management of insect pests with Bt crops in the United States. *Annu. Rev. Entomol.* **68**, 31–49 (2023). doi: [10.1146/annurev-ento-102020-105502](https://doi.org/10.1146/annurev-ento-102020-105502); pmid: [36170641](https://pubmed.ncbi.nlm.nih.gov/36170641/)
- V. V. Krishna, M. Qaim, Bt cotton and sustainability of pesticide reductions in India. *Agric. Syst.* **107**, 47–55 (2012). doi: [10.1016/j.agsy.2011.11.005](https://doi.org/10.1016/j.agsy.2011.11.005)
- K. R. Kranthi, G. D. Stone, Long-term impacts of Bt cotton in India. *Nat. Plants* **6**, 188–196 (2020). doi: [10.1038/s41477-020-0615-5](https://doi.org/10.1038/s41477-020-0615-5); pmid: [32170289](https://pubmed.ncbi.nlm.nih.gov/32170289/)
- Y. Carrière *et al.*, “Crop rotation mitigates impacts of corn rootworm resistance to transgenic Bt corn” in *Proceedings of the National Academy of Sciences of the United States* (PNAS, 2019), pp. 18385–18392. doi: [10.1073/pnas.2003604117](https://doi.org/10.1073/pnas.2003604117)
- P. Bagla, Hardy cotton-munching pests are latest blow to GM crops. *Science* **327**, 1439 (2010). doi: [10.1126/science.1190242](https://doi.org/10.1126/science.1190242)
- A. J. Gassmann, J. L. Petzold-Maxwell, R. S. Keweshan, M. W. Dunbar, Field-evolved resistance to Bt maize by western corn rootworm. *PLOS ONE* **6**, e22629 (2011). doi: [10.1371/journal.pone.0022629](https://doi.org/10.1371/journal.pone.0022629); pmid: [21829470](https://pubmed.ncbi.nlm.nih.gov/21829470/)
- B. E. Tabashnik, J. A. Fabrick, Y. Carrière, Global patterns of insect resistance to transgenic Bt crops: The first 25 years. *J. Econ. Entomol.* **116**, 297–309 (2023). doi: [10.1093/jeet/toac183](https://doi.org/10.1093/jeet/toac183); pmid: [36610076](https://pubmed.ncbi.nlm.nih.gov/36610076/)



64. Y. Carrière *et al.*, Crop rotation mitigates impacts of corn rootworm resistance to transgenic Bt corn. *Proc. Natl. Acad. Sci. U.S.A.* **117**, 18385–18392 (2020). doi: [10.1073/pnas.2003604117](https://doi.org/10.1073/pnas.2003604117); pmid: [32690686](https://pubmed.ncbi.nlm.nih.gov/32690686/)
65. M. Qaim, G. Traxler, Roundup Ready soybeans in Argentina: Farm level and aggregate welfare effects. *Agric. Econ.* **32**, 73–86 (2005). doi: [10.1111/j.0169-5150.2005.00006.x](https://doi.org/10.1111/j.0169-5150.2005.00006.x)
66. C. M. Benbrook, Trends in glyphosate herbicide use in the United States and globally. *Environ. Sci. Eur.* **28**, 3 (2016). doi: [10.1186/s12302-016-0070-0](https://doi.org/10.1186/s12302-016-0070-0); pmid: [27752438](https://pubmed.ncbi.nlm.nih.gov/27752438/)
67. B. G. Silva Pinto, T. K. Marques Soares, M. Azevedo Linhares, N. Castilhos Ghisi, Occupational exposure to pesticides: Genetic danger to farmworkers and manufacturing workers - A meta-analytical review. *Sci. Total Environ.* **748**, 141382 (2020). doi: [10.1016/j.scitotenv.2020.141382](https://doi.org/10.1016/j.scitotenv.2020.141382); pmid: [32818891](https://pubmed.ncbi.nlm.nih.gov/32818891/)
68. C. E. Handford, C. T. Elliott, K. Campbell, A review of the global pesticide legislation and the scale of challenge in reaching the global harmonization of food safety standards. *Integr. Environ. Assess. Manag.* **11**, 525–536 (2015). doi: [10.1002/ieam.1635](https://doi.org/10.1002/ieam.1635); pmid: [25765969](https://pubmed.ncbi.nlm.nih.gov/25765969/)
69. S. J. Smyth, The human health benefits from GM crops. *Plant Biotechnol. J.* **18**, 887–888 (2020). doi: [10.1111/pbi.13261](https://doi.org/10.1111/pbi.13261); pmid: [31544299](https://pubmed.ncbi.nlm.nih.gov/31544299/)
70. S. Kouser, M. Qaim, Impact of Bt cotton on pesticide poisoning in smallholder agriculture: A panel data analysis. *Ecol. Econ.* **70**, 2105–2113 (2011). doi: [10.1016/j.ecolecon.2011.06.008](https://doi.org/10.1016/j.ecolecon.2011.06.008)
71. G. C. Nelson, D. S. Bullock, Simulating a relative environmental effect of glyphosate-resistant soybeans. *Ecological Economics* **45**, 189–202 (2003). doi: [10.1016/j.ecolecon.2011.06.008](https://doi.org/10.1016/j.ecolecon.2011.06.008)
72. IARC, *Some Organophosphate Insecticides and Herbicides*. (WHO, 2024). Vol. 112.
73. M. Desquilbet, D. S. Bullock, F. M. D'Arcangelo, A discussion of the market and policy failures associated with the adoption of herbicide-tolerant crops. *International Journal of Agricultural Sustainability* **17**, 326–337, (2019). doi: [10.1016/j.ecolecon.2011.06.008](https://doi.org/10.1016/j.ecolecon.2011.06.008)
74. N. Defarge *et al.*, Co-formulants in glyphosate-based herbicides disrupt aromatase activity in human cells below toxic levels. *Int. J. Environ. Res. Public Health* **13**, 264 (2016). doi: [10.3390/ijerph13030264](https://doi.org/10.3390/ijerph13030264); pmid: [26927151](https://pubmed.ncbi.nlm.nih.gov/26927151/)
75. I. M. Meftaul *et al.*, Controversies over human health and ecological impacts of glyphosate: Is it to be banned in modern agriculture? *Environ. Pollut.* **263**, 114372 (2020). doi: [10.1016/j.envpol.2020.114372](https://doi.org/10.1016/j.envpol.2020.114372); pmid: [32203845](https://pubmed.ncbi.nlm.nih.gov/32203845/)
76. E. D. Perry, G. Moschini, Neonicotinoids in US maize: Insecticide substitution effects and environmental risk. *J. Environ. Econ. Manage.* **102**, 102320 (2020). doi: [10.1016/j.jeem.2020.102320](https://doi.org/10.1016/j.jeem.2020.102320)
77. A. R. Kniss, Long-term trends in the intensity and relative toxicity of herbicide use. *Nat. Commun.* **8**, 14865 (2017). doi: [10.1038/ncomms14865](https://doi.org/10.1038/ncomms14865); pmid: [28393866](https://pubmed.ncbi.nlm.nih.gov/28393866/)
78. S. Lee, G. Moschini, E. D. Perry, Genetically engineered varieties and applied pesticide toxicity in US maize and soybeans: Heterogeneous and evolving impacts. *Ecol. Econ.* **211**, 107873 (2023). doi: [10.1016/j.ecolecon.2023.107873](https://doi.org/10.1016/j.ecolecon.2023.107873)
79. B. K. Armstrong *et al.*, Differences in the carcinogenic evaluation of glyphosate between the International agency for research on cancer (IARC) and the European food safety authority (EFSA). *J. Epidemiol. Community Health* **70**, 741–745 (2016). doi: [10.1016/j.ecolecon.2023.107873](https://doi.org/10.1016/j.ecolecon.2023.107873); pmid: [26941213](https://pubmed.ncbi.nlm.nih.gov/26941213/)
80. C. J. Portier, A comprehensive analysis of the animal carcinogenicity data for glyphosate from chronic exposure rodent carcinogenicity studies. *Environmental Health: A Global Access Science Source* **19**, 1–18 (2020). doi: [10.1016/j.ecolecon.2023.107873](https://doi.org/10.1016/j.ecolecon.2023.107873)
81. C. M. Benbrook, Why Regulators Lost Track and Control of Pesticide Risks: Lessons From the Case of Glyphosate-Based Herbicides and Genetically Engineered-Crop Technology. *Curr. Environ. Health Rep.* **5**, 387–395 (2018). doi: [10.1007/s40572-018-0207-y](https://doi.org/10.1007/s40572-018-0207-y); pmid: [30003510](https://pubmed.ncbi.nlm.nih.gov/30003510/)
82. T. R. Stafford, L. B. Best, Bird response to grit and pesticide granule characteristics: Implications for risk assessment and risk reduction. *Environ. Toxicol. Chem.* **18**, 722–733 (2009). doi: [10.1002/etc.5620180419](https://doi.org/10.1002/etc.5620180419)
83. D. R. J. Moore, D. L. Fischer, R. S. Teed, S. I. Rodney, Probabilistic risk-assessment model for birds exposed to granular pesticides. *Integr. Environ. Assess. Manag.* **6**, 260–272 (2010). doi: [10.1897/IEAM\\_2009-021.1](https://doi.org/10.1897/IEAM_2009-021.1); pmid: [20821688](https://pubmed.ncbi.nlm.nih.gov/20821688/)
84. M. E. Skidmore, K. M. Sims, H. K. Gibbs, Agricultural intensification and childhood cancer in Brazil. *Proc. Natl. Acad. Sci. U.S.A.* **120**, e2306003120 (2023). doi: [10.1073/pnas.2306003120](https://doi.org/10.1073/pnas.2306003120); pmid: [37903255](https://pubmed.ncbi.nlm.nih.gov/37903255/)
85. S. Sankula, G. Marmon, E. Blumenthal, "Biotechnology-derived crops planted in 2004: Impacts on US agriculture" (National Center for Food and Agricultural Policy, 2005).
86. J. Fernandez-Cornejo, C. Hallahan, R. F. Nehring, S. Wechsler, A. Grube, Conservation tillage, herbicide use, and genetically engineered crops in the United States: The case of soybeans. *AgBioForum* **15**, 231–241 (2013).
87. W. A. Givens *et al.*, Survey of tillage trends following the adoption of glyphosate-resistant crops. *Weed Technol.* **23**, 150–155 (2017). doi: [10.1614/WT-08-038.1](https://doi.org/10.1614/WT-08-038.1)
88. E. Perry, G. Moschini, D. A. Hennessy, Testing for complementarity: Glyphosate tolerant soybeans and conservation tillage. *Am. J. Agric. Econ.* **98**, 765–784 (2016). doi: [10.1093/ajae/aaw001](https://doi.org/10.1093/ajae/aaw001)
89. C. M. Pittelkow *et al.*, Productivity limits and potentials of the principles of conservation agriculture. *Nature* **517**, 365–368 (2015). doi: [10.1038/nature13809](https://doi.org/10.1038/nature13809); pmid: [25337882](https://pubmed.ncbi.nlm.nih.gov/25337882/)
90. J. M. Holland, The environmental consequences of adopting conservation tillage in Europe: Reviewing the evidence. *Agric. Ecosyst. Environ.* **103**, 1–25 (2004). doi: [10.1016/j.jagee.2003.12.018](https://doi.org/10.1016/j.jagee.2003.12.018)
91. A. P. Behrer, D. Lobell, Higher levels of no-till agriculture associated with lower PM<sub>2.5</sub> in the Corn Belt. *Environ. Res. Lett.* **17**, 094012 (2022). doi: [10.1088/1748-9326/ac816f](https://doi.org/10.1088/1748-9326/ac816f)
92. F. Cozím-Melges *et al.*, Farming practices to enhance biodiversity across biomes: A systematic review. *npj Biodiversity* **3**, 1 (2024). doi: [10.1038/s4185-023-00034-2](https://doi.org/10.1038/s4185-023-00034-2)
93. J. M. Baker, T. E. Ochsner, R. T. Venterea, T. J. Griffis, Tillage and soil carbon sequestration—What do we really know? *Agric. Ecosyst. Environ.* **118**, 1–5 (2007). doi: [10.1016/j.jagee.2006.05.014](https://doi.org/10.1016/j.jagee.2006.05.014)
94. R. S. Nicoloso, C. W. Rice, Intensification of no-till agricultural systems: An opportunity for carbon sequestration. *Soil Sci. Soc. Am. J.* **85**, 1395–1409 (2021). doi: [10.1002/saj2.20260](https://doi.org/10.1002/saj2.20260)
95. J.-P. Chavas, G. Shi, J. Lauer, The effects of GM technology on maize yield. *Crop Sci.* **54**, 1331–1335 (2014). doi: [10.2135/cropsci2013.10.0709](https://doi.org/10.2135/cropsci2013.10.0709)
96. E. Strobl, Preserving local biodiversity through crop diversification. *Am. J. Agric. Econ.* **104**, 1140–1174 (2022). doi: [10.1111/ajae.12265](https://doi.org/10.1111/ajae.12265)
97. A. E. Larsen, F. Noack, Identifying the landscape drivers of agricultural insecticide use leveraging evidence from 100,000 fields. *Proc. Natl. Acad. Sci. U.S.A.* **114**, 5473–5478 (2017). doi: [10.1073/pnas.1620674114](https://doi.org/10.1073/pnas.1620674114); pmid: [28484021](https://pubmed.ncbi.nlm.nih.gov/28484021/)
98. T. M. Bowles *et al.*, Long-term evidence shows that crop-rotation diversification increases agricultural resilience to adverse growing conditions in North America. *One Earth* **2**, 284–293 (2020). doi: [10.1016/j.oneear.2020.02.007](https://doi.org/10.1016/j.oneear.2020.02.007)
99. K. Emerick, A. De Janvry, E. Sadoulet, M. H. Dar, Technological innovations, downside risk, and the modernization of agriculture. *Am. Econ. Rev.* **106**, 1537–1561 (2016). doi: [10.1257/aer.20150474](https://doi.org/10.1257/aer.20150474)
100. C. A. Taylor, G. Heal, "Fertilizer and Algal Blooms" in *Risks in Agricultural Supply Chains*, P. Antràs, D. Zilberman, Eds. (Univ. of Chicago Press, 2023).
101. M. Cuhra, Review of GMO safety assessment studies: Glyphosate residues in Roundup Ready crops is an ignored issue. *Environ. Sci. Eur.* **27**, 20 (2015). doi: [10.1186/s12302-015-0052-7](https://doi.org/10.1186/s12302-015-0052-7)
102. D. A. Andow, C. Zwhalen, Assessing environmental risks of transgenic plants. *Ecol. Lett.* **9**, 196–214 (2006). doi: [10.1111/j.1461-0248.2005.00846.x](https://doi.org/10.1111/j.1461-0248.2005.00846.x); pmid: [16958885](https://pubmed.ncbi.nlm.nih.gov/16958885/)
103. N. Defarge, M. Otto, A. Hilbeck, A roundup herbicide causes high mortality and impairs development of chrysoperla carnea (stephens) (neuroptera: chrysopidae). *SSRN Electronic Journal* **11** (2023).doi: [10.1016/j.scitotenv.2022.161158](https://doi.org/10.1016/j.scitotenv.2022.161158)
104. H. J. Venter, T. Böhn, Interactions between Bt crops and aquatic ecosystems: A review. *Environ. Toxicol. Chem.* **35**, 2891–2902 (2016). doi: [10.1002/etc.3583](https://doi.org/10.1002/etc.3583); pmid: [27530353](https://pubmed.ncbi.nlm.nih.gov/27530353/)
105. A. Pott, M. Otto, R. Schulz, Impact of genetically modified organisms on aquatic environments: Review of available data for the risk assessment. *Sci. Total Environ.* **635**, 687–698 (2018). doi: [10.1016/j.scitotenv.2018.04.013](https://doi.org/10.1016/j.scitotenv.2018.04.013); pmid: [29680759](https://pubmed.ncbi.nlm.nih.gov/29680759/)
106. N. C. Ellstrand, When transgenes wander, should we worry? *Plant Physiol.* **125**, 1543–1545 (2001). doi: [10.1104/pp.125.4.1543](https://doi.org/10.1104/pp.125.4.1543); pmid: [11299333](https://pubmed.ncbi.nlm.nih.gov/11299333/)
107. D. A. Levin, J. Francisco-Ortega, R. K. Jansen, Hybridization and the extinction of rare plant species. *Conserv. Biol.* **10**, 10–16 (1996). doi: [10.1046/j.1523-1739.1996.10010010.x](https://doi.org/10.1046/j.1523-1739.1996.10010010.x)
108. N. Ellstrand, P. Meirmans, J. Rong, K. Vrieling, D. Hooftman, Introgression of Crop Alleles into Wild or Weedy Populations. *Annu. Rev. Ecol. Evol. Syst.* **44**, 325–345 (2013). doi: [10.1146/annurev-ecolsys-110512-135840](https://doi.org/10.1146/annurev-ecolsys-110512-135840)
109. B.-R. Lu, X. Yang, N. C. Ellstrand, Fitness correlates of crop transgene flow into weed populations: A case study of weedy rice in China and other examples. *Evol. Appl.* **9**, 857–870 (2016). doi: [10.1111/eva.12377](https://doi.org/10.1111/eva.12377); pmid: [27468304](https://pubmed.ncbi.nlm.nih.gov/27468304/)
110. N. C. Ellstrand, L. H. Rieseberg, When gene flow really matters: Gene flow in applied evolutionary biology. *Evol. Appl.* **9**, 833–836 (2016). doi: [10.1111/eva.12402](https://doi.org/10.1111/eva.12402); pmid: [27468301](https://pubmed.ncbi.nlm.nih.gov/27468301/)
111. M. A. Tsiafouli *et al.*, Intensive agriculture reduces soil biodiversity across Europe. *Glob. Change Biol.* **21**, 973–985 (2015). doi: [10.1111/gcb.12752](https://doi.org/10.1111/gcb.12752); pmid: [25242445](https://pubmed.ncbi.nlm.nih.gov/25242445/)
112. F. H. Tang, M. Lenzen, A. McBratney, F. Maggi, Risk of pesticide pollution at the global scale. *Nat. Geosci.* **14**, 206–210 (2021). doi: [10.1038/s41561-021-00712-5](https://doi.org/10.1038/s41561-021-00712-5)
113. J. Candel, G. Pe'er, R. Finger, Science calls for ambitious European pesticide policies. *Nat. Food* **4**, 272–272 (2023). doi: [10.1038/s43016-023-00727-8](https://doi.org/10.1038/s43016-023-00727-8); pmid: [37117539](https://pubmed.ncbi.nlm.nih.gov/37117539/)
114. Y. Li, R. Miao, M. Khanna, Neonicotinoids and decline in bird biodiversity in the United States. *Nat. Sustain.* **3**, 1027–1035 (2020). doi: [10.1038/s41893-020-0582-x](https://doi.org/10.1038/s41893-020-0582-x)
115. B. Strandberg *et al.*, Effects of glyphosate spray-drift on plant flowering. *Environ. Pollut.* **280**, 116953 (2021). doi: [10.1016/j.envpol.2021.116953](https://doi.org/10.1016/j.envpol.2021.116953); pmid: [33784566](https://pubmed.ncbi.nlm.nih.gov/33784566/)
116. Y. L. Dupont, B. Strandberg, C. Damgaard, Effects of herbicide and nitrogen fertilizer on non-target plant reproduction and indirect effects on pollination in *Tanacetum vulgare* (Asteraceae). *Agric. Ecosyst. Environ.* **262**, 76–82 (2018). doi: [10.1016/j.jagee.2018.04.014](https://doi.org/10.1016/j.jagee.2018.04.014)
117. C. Sirami *et al.*, Increasing crop heterogeneity enhances multitrophic diversity across agricultural regions. *Proc. Natl. Acad. Sci. U.S.A.* **116**, 16442–16447 (2019). doi: [10.1073/pnas.1906419116](https://doi.org/10.1073/pnas.1906419116); pmid: [31358630](https://pubmed.ncbi.nlm.nih.gov/31358630/)
118. L. Kehoe *et al.*, Biodiversity at risk under future cropland expansion and intensification. *Nat. Ecol. Evol.* **1**, 1129–1135 (2017). doi: [10.1038/s41559-017-0234-3](https://doi.org/10.1038/s41559-017-0234-3); pmid: [29046577](https://pubmed.ncbi.nlm.nih.gov/29046577/)
119. D. Tilman *et al.*, Future threats to biodiversity and pathways for their prevention. *Nature* **546**, 73–81 (2017). doi: [10.1038/nature22900](https://doi.org/10.1038/nature22900); pmid: [28569796](https://pubmed.ncbi.nlm.nih.gov/28569796/)
120. E. V. S. Motta, K. Raymann, N. A. Moran, Glyphosate perturbs the gut microbiota of honey bees. *Proc. Natl. Acad. Sci. U.S.A.* **115**, 10305–10310 (2018). doi: [10.1073/pnas.1803880115](https://doi.org/10.1073/pnas.1803880115); pmid: [30249635](https://pubmed.ncbi.nlm.nih.gov/30249635/)
121. M.-P. Hébert *et al.*, Widespread agrochemicals differentially affect zooplankton biomass and community structure. *Ecol. Appl.* **31**, e02423 (2021). doi: [10.1002/eap.2423](https://doi.org/10.1002/eap.2423); pmid: [34288209](https://pubmed.ncbi.nlm.nih.gov/34288209/)
122. A. Weidenmüller, A. Meltzer, S. Neupert, A. Schwarz, C. Kleinedam, Glyphosate impairs collective thermoregulation in bumblebees. *Science* **376**, 1122–1126 (2022). doi: [10.1126/science.abf7482](https://doi.org/10.1126/science.abf7482); pmid: [35653462](https://pubmed.ncbi.nlm.nih.gov/35653462/)
123. S. Klatiyk *et al.*, Aquatic ecotoxicity of glyphosate, its formulations, and co-formulants: Evidence from 2010 to 2023. *Environ. Sci. Eur.* **36**, 22 (2024). doi: [10.1186/s12302-024-00849-1](https://doi.org/10.1186/s12302-024-00849-1)
124. D. Da Mata, M. Dotta, T. Lobo, "Technological Progress and Climate Change: Evidence from the Agricultural Sector" (SSRN, 2023); [https://papers.ssrn.com/sol3/papers.cfm?abstract\\_id=4358828](https://papers.ssrn.com/sol3/papers.cfm?abstract_id=4358828)
125. M. Dainese *et al.*, A global synthesis reveals biodiversity-mediated benefits for crop production. *Sci. Adv.* **5**, eaax0121 (2019). doi: [10.1126/sciadv.aax0121](https://doi.org/10.1126/sciadv.aax0121); pmid: [31663019](https://pubmed.ncbi.nlm.nih.gov/31663019/)
126. G. Tamburini *et al.*, Agricultural diversification promotes multiple ecosystem services without compromising yield. *Sci. Adv.* **6**, eaba1715 (2020). doi: [10.1126/sciadv.aax0121](https://doi.org/10.1126/sciadv.aax0121); pmid: [31663019](https://pubmed.ncbi.nlm.nih.gov/31663019/)

127. D. Beillouin, T. Ben-Ari, E. Malézieux, V. Seufert, D. Makowski, Positive but variable effects of crop diversification on biodiversity and ecosystem services. *Glob. Change Biol.* **27**, 4697–4710 (2021). doi: [10.1111/gcb.15747](https://doi.org/10.1111/gcb.15747); pmid: [34114719](https://pubmed.ncbi.nlm.nih.gov/34114719/)

128. L. C. Ponisio *et al.*, Diversification practices reduce organic to conventional yield gap. *Proc. Biol. Sci.* **282**, 20141396 (2015). doi: [10.1098/rspb.2014.1396](https://doi.org/10.1098/rspb.2014.1396); pmid: [25621333](https://pubmed.ncbi.nlm.nih.gov/25621333/)

129. Council for Agricultural Science and Technology (CAST), “Applications, Benefits, and Challenges of Genome Edited Crops” (CAST technical report no. 74, 2024).

130. Food and Agriculture Organization of the United States (FAO), “FAOSTAT” (FAO, 2024); <https://www.fao.org/faostat/en/#data>.

ACKNOWLEDGMENTS

We thank the two anonymous reviewers, and S. Heredia for her excellent help with the overview figure. **Funding:** F.N. acknowledges support from the Canada Research Chairs Program (Canada Research Chair in Economic and Environmental Interactions). F.N., R.S., and E.S.-R. acknowledge support from the SSHRC Insight Grant (AWD-021156). R.S. and F.N. acknowledge funding from UBC’s Faculty of Land and Food Systems Internal Research Grant and Grants for Catalyzing Research Clusters to UBC’s Biodiversity Research Centre. M.Q. acknowledges support from Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) (EXC-2070-390732324-PhenoRob). A.L. acknowledges support from the Agriculture and Food Research Initiative of the National Institute of Food and

Agriculture, USDA (2022-67019-36397, 2023-68016-40683) and the US National Science Foundation (2042526). L.M.G. acknowledges funding from NSERC (Natural Sciences and Engineering Research Council of Canada) Discovery Grant. **Competing interests:** All authors declare no competing interests. **License information:** Copyright © 2024 the authors, some rights reserved; exclusive licensee American Association for the Advancement of Science. No claim to original US government works. <https://www.science.org/about/science-licenses-journal-article-reuse>

Submitted 27 February 2024; accepted 24 June 2024  
10.1126/science.ado9340